


CHAPTER 4 A Tour of the Cell

YOU MUST KNOW


- Three differences between prokaryotic and eukaryotic cells.
- The structure and function of organelles common to plant and animal cells.
- The structure and function of organelles found only in plant cells or only in animal cells.
- How different cell types show differences in subcellular components.
- How internal membranes and organelles contribute to cell functions.
- How cell size and shape affect the overall rate of nutrient intake and waste elimination.

2 TYPES OF CELLS:

- 1. <u>Prokaryotes</u>: Domain Bacteria & Archaea
- 2. <u>Eukaryotes</u> (Domain Eukarya): Protists, Fungi, Plants, Animals




Animal Cells



Human cells from lining of uterus (colorized TEM)

© 2011 Pearson Education, In-

A PROKARYOTIC CELL (BACTERIA)



PROKARYOTE VS. EUKARYOTE

- "before" "kernel"
- No nucleus
- DNA in a nucleoid
- Cytosol
- No organelles other than ribosomes
- Small size
- Primitive
- o i.e. Bacteria & Archaea


- "true" "kernel"
- Has nucleus and nuclear envelope
- Cytosol
- Membrane-bound organelles with specialized structure/ function
- Much larger in size
- More complex
- i.e. plant/animal cell

CELL SIZE AND SCALE


http://learn.genetics.utah.edu/content/begin/cells/scale/

Scale of the Universe:

http://www.onemorelevel.com/game/scale_of_the_universe_2012




- Cells must be small to maintain a large surface area to volume ratio
- Large S.A. allows ↑ rates of chemical exchange between cell and environment




CALCULATE VOLUME OF A SPHERE:

$$V=\frac{4}{3}\pi r^3$$



© 2016 Pearson Education, Inc.


SURFACE AREA EXAMPLE (ANIMAL):




Small Intestine: highly folded surface to increase absorption of nutrients

- <u>Villi</u>: finger-like projections on SI wall
- Microvilli: projections on each cell

Folds -> Villi -> Microvill OF SMALL INTESTINE






© 2011 Pearson Education, Inc.

SURFACE AREA EXAMPLE (PLANT):


Root hairs: extensions of root epidermal cells; increase surface area for absorbing water and minerals






NUCLEUS


- <u>Function</u>: control center of cell
- Contains DNA (& mRNA)
- Surrounded by double membrane (nuclear envelope)
 - Continuous with the rough ER
- Nuclear pores: control what enters/leaves nucleus
- <u>Chromatin</u>: complex of DNA + proteins; makes up chromosomes
- <u>Nucleolus</u>: region where ribosomal subunits (rRNA + proteins) are formed






RIBOSOMES

- Function: protein synthesis
- Composed of rRNA + protein
- Large subunit + small subunit
- Types:
 - 1. <u>Free ribosomes</u>: float in cytosol, produce proteins used within cell
 - 2. <u>Bound ribosomes</u>: attached to ER, make proteins for export from cell






SYSTEM:

Regulates protein traffic & performs metabolic functions

ENDOPLASMIC RETICULUM (ER)


- Network of membranes and sacs
- Types:
 - 1. Rough ER: ribosomes on surface
 - <u>Function</u>: package proteins for secretion, send transport vesicles to Golgi, make replacement membrane
 - 2. Smooth ER: no ribosomes on surface
 - <u>Function</u>: synthesize lipids, metabolize carbs, detox drugs & poisons, store Ca²⁺

ENDOPLASMIC RETICULUM (ER)




GOLGI APPARATUS


- <u>Function</u>: synthesis & packaging of materials (small molecules) for transport (in vesicles); produce lysosomes
- Series of flattened membrane sacs (cisternae)
 - Cis face: receives vesicles
 - *Trans* face: ships vesicles





LYSOSOMES

- <u>Function</u>: intracellular digestion; recycle cell's materials; programmed cell death (apoptosis)
- Contains hydrolytic enzymes











(a) Phagocytosis


(b) Autophagy


VACUOLES


- <u>Function</u>: storage of materials (food, water, minerals, pigments, poisons)
- Membrane-bound vesicles
- Eg. food vacuoles, contractile vacuoles
- Plants: <u>large central vacuole</u> -- stores water, ions; retains water for turgor pressure















MITOCHONDRIA

- <u>Function</u>: site of cellular respiration
- Double membrane: outer and inner membrane
- Cristae: folds of inner membrane; contains enzymes for ATP production; increased surface area to ↑ ATP made
- Matrix: fluid-filled inner compartment



(b) Network of mitochondria in a protist cell (LM)

Nuclear DNA

Mitochondrial


DNA


Mitochondria

(a) Diagram and TEM of mitochondrion

CHLOROPLASTS

- <u>Function</u>: site of photosynthesis
- Double membrane
- Thylakoid disks in stacks (grana); stroma (fluid)
- Contains chlorophylls (pigments) for capturing sunlight energy






(b) Chloroplasts in an algal cell

© 2011 Pearson Education, Inc.


ENDOSYMBIONT THEORY



- Mitochondria & chloroplasts share similar origin
- Prokaryotic cells engulfed by ancestors of eukaryotic cells
 - <u>Evidence:</u>
 - Double-membrane structure
 - Have own ribosomes & DNA
 - Reproduce independently within cell


PEROXISOMES

- <u>Functions</u>: break down fatty acids; detox alcohol
- Involves production of hydrogen peroxide (H₂O₂)



CYTOSKELETON: NETWORK OF PROTEIN FIBERS

• <u>Function</u>: support, motility, regulate biochemical activities




10 mm



Table 4.1 The Structure and Function of the Cytoskeleton				
Property	Microtubules (Tubulin Polymers)	Microfilaments (Actin Filaments)	Intermediate Filaments	
Structure	Hollow tubes	Two intertwined strands of actin	Fibrous proteins coiled into cables	
Diameter	25 nm with 15-nm lumen	7 nm	8–12 nm	
Protein subunits	Tubulin, a dimer consisting of α-tubulin and β-tubulin	Actin	One of several different proteins (such as keratins)	
Main functions	Maintenance of cell shape; cell mo- tility; chromosome movements in cell division; organelle movements	Maintenance of cell shape; changes in cell shape; muscle contraction; cytoplasmic streaming (plant cells); cell motility; cell division (animal cells)	Maintenance of cell shape; anchorage of nucleus and certain other organelles; formation of nuclear lamina	
Fluorescence micrographs of fibroblasts. Fibroblasts are a favorite cell type for cell biology studies because they spread out flat and their internal structures are easy to see. In each, the structure of interest has been tagged with fluorescent molecules. The DNA in the nucleus has also been tagged in the first micrograph (blue) and third micrograph (orange).	Column of tubulin dimers 25 nm γ Tubulin dimer	Actin subunit 7 nm	Keratin proteins Fibrous subunit (keratins coiled together) 8–12 nm	


PLANT CELLS

- <u>Cell wall</u>: protect plant, maintain shape
 - Composed of cellulose
- Plasmodesmata:
 channels between cells to
 allow passage of molecules
 from cell to cell



EXTRACELLULAR MATRIX (ECM)


- Outside plasma membrane of animal cells
- Composed of glycoproteins (ex. collagen)
- <u>Function</u>: Strengthens tissues and transmits external signals to cell



INTERCELLULAR JUNCTIONS (ANIMAL CELLS)

- <u>Tight junctions</u>: 2 cells are fused to form watertight seal
- Desmosomes: "rivets" that fasten adjacent cells into strong sheets
- Gap junctions:

 channels through which ions, sugar, small molecules can pass



Plant Cells Only	Animals Cells Only	
Central vacuoles	Lysosomes	
Chloroplasts	Centrioles	
Cell wall of cellulose	Flagella, cilia	
Plasmodesmata	Desmosomes, tight and gap junctions	
	Extracellular matrix (ECM)	

HARVARD CELL VIDEO


http://multimedia.mcb.harvard.edu/ anim innerlife.html

Cell Component	Structure	Function
Nucleus (ER)	Surrounded by nuclear envelope (double membrane) perforated by nuclear pores; nuclear envelope continuous with endoplasmic reticulum (ER)	Houses chromosomes, which are made of chromatin (DNA and proteins); contains nucleoli, where ribosomal subunits are made; pores regulate entry and exit of materials
Ribosome	Two subunits made of ribosomal RNA and proteins; can be free in cytosol or bound to ER	Protein synthesis

© 2016 Pearson Education, Inc.

Cell Component	Structure	Function
Endoplasmic reticulum (Nuclear envelope)	Extensive network of membrane-bounded tubules and sacs; membrane separates lumen from cytosol; continuous with nuclear envelope	Smooth ER: synthesis of lipids, metabolism of carbohydrates, Ca ²⁺ storage, detoxification of drugs and poisons Rough ER: aids in synthesis of secretory and other proteins from bound ribosomes; adds carbohydrates to proteins to make glycoproteins; produces new membrane
Golgi apparatus	Stacks of flattened membranous sacs; has polarity (<i>cis</i> and <i>trans</i> faces)	Modification of proteins, carbohydrates on proteins, and phospholipids; synthesis of many polysaccharides; sorting of Golgi products, which are then released in vesicles
Lysosome	Membranous sac of hydrolytic enzymes (in animal cells)	Breakdown of ingested substances, cell macromolecules, and damaged organelles for recycling
Vacuole	Large membrane-bounded vesicle	Digestion, storage, waste disposal, water balance, plant cell growth and protection

Cell Component	Structure	Function
Mitochondrion	Bounded by double membrane; inner membrane has infoldings (cristae)	Cellular respiration
Chloroplast	Typically two membranes around fluid stroma, which contains thylakoids stacked into grana (in cells of photosynthetic eukaryotes, including plants)	Photosynthesis
Peroxisome	Specialized metabolic compartment bounded by a single membrane	Contains enzymes that transfer hydrogen atoms from certain molecules to oxygen, producing hydrogen peroxide (H ₂ O ₂) as a by-product; H ₂ O ₂ is converted to water by another enzyme




Smooth Endoplasmic Reticulum


might-ochondria



definitely-chondria







mitochondria




-tumblr famous -only drinks gatorade -powerhouse of the cell

nucleus



-control freak
-"everyone is important uwu"
-unofficial squad leader

ribosome




-actually works for a living -always making things for their friends -highkey wants to be nucleus's bff

nucleolus




-nucleus's actual bff -superiority complex -basically 12 years old

golgi apparatus



-loves duct tape -gives great advice -unproblematic fave

lysosome



-always salty -loves dogs -will fight anyone

ER



-total mom friend -better than you at literally everything -only one who can drive

vacuole



-really quiet except when they're not -v dependable -does crossfit

chloroplast



-super elitist
-only wears green except
on st patricks day
-lowkey hippie