General Chemistry Chapter 5 Pre-Test

- 1.) (2 pts each, 16 pts total) Use your knowledge of ideal gas laws to answer each of the following. Assume all other relevant factors are constant.
 - a) As the pressure of an ideal gas increases, the volume must (increase or decrease)
 - b) As the volume of an ideal gas decreases, the temperature must (increase or decrease) PV = nRT
 - c) As the volume of an ideal gas decreases, the pressure must (increase or decrease)
 - d) As the temperature of an ideal gas increases, the pressure must increase or decrease)
 - e) As the amount of an ideal gas increases the pressure must (increase or decrease)
 - f) As the amount of an ideal gas decreases, the volume must (increase or decrease)
 - g) As the pressure of an ideal gas increases, the temperature must (increase or decrease)
 - h) As the temperature of an ideal gas increases, the volume must (increase or decrease)

2.) (12 pts) A sample of argon gas has a pressure of 628 torr and a volume of 2.54 L. What is the volume of the gas if the pressure is adjusted to 846 torr?

$$P_{1}V_{1} = P_{2}V_{2}$$
 $P_{1}V_{1} = P_{2}V_{2}$
 $P_{2}V_{1} = P_{2}V_{2}$
 $P_{3}V_{2} = P_{3}V_{2}$
 $P_{4}V_{1} = P_{2}V_{2}$
 $P_{5}V_{1} = P_{5}V_{2}$
 $P_{5}V_{2} = P_{5}V_{1}$
 $P_{5}V_{1} = P_{5}V_{2}$
 $P_{5}V_{2} = P_{5}V_{1}$
 $P_{5}V_{2} = P_{5}V_{1}$
 $P_{5}V_{1} = P_{5}V_{2}$
 $P_{5}V_{5} = P_{5}V_{5}$
 $P_{5}V_{5} = P_{5}V_{5}$

3.) (12 pts) A 7.84 L sample of carbon dioxide gas has a temperature of 38.0 °C. What is the volume of the same gas if the temperature is adjusted to 52.0 °C?

$$\frac{V_{1}}{T_{1}} = \frac{V_{2}}{V_{2}} \quad V_{1} = 7.84 \qquad PV = nRT \\
T_{1} \quad T_{2} \quad T_{1} = 38 + 273 \qquad P \qquad P \qquad P$$

$$\frac{311k}{V_{2}} = 7 \quad T_{2} = 52 + 273 \qquad T \qquad V = \frac{nRT}{P} / 7$$

$$\frac{325 k}{T_{1}} \quad T_{2} \quad V_{2} \quad T_{2} \qquad V_{2} = \frac{T_{2} V_{1}}{T_{1}} = \frac{(325k)(7.84L)}{311k}$$

$$\frac{311k}{8.19L}$$

4.) (12 pts) A sealed container of water vapor has a pressure of 1.00 atm and a temperature of 37 °C. What is the temperature of the sealed gas if the pressure is increased to 1.86 atm?

$$P_{1} = 1.00 \text{ atm} \quad T_{1} = 37 + 273 = 310 \text{ K}$$

$$P_{2} = 1.86 \text{ atm} \quad T_{2} = ?$$

$$P_{1} \times \frac{P_{2}}{T_{2}} = \frac{1}{V} =$$

5.) (12 pts) What is the volume of a 0.74 mol sample of oxygen gas at 0.88 atm at a temperature of 78.0 °C?

$$PV = nRT \qquad n = 0.74 \text{ mol } P = 0.88 \text{ atm}$$

$$T = 78 + 273 = 351K$$

$$V = nRT = (0.74 \text{ mol})(0.0921)(351K)$$

$$0.98 \text{ atm}$$

$$24.2L$$

6.) (12 pts) A 1.85 L container of 4.92 g of an unknown ideal gas is measured at 1.50 atm and 29.0 °C. What is the molar mass of the gas?

reaction prior to solving.

$$\frac{1}{2}$$

$$V = \frac{NRT}{P} = \frac{(0.5 \text{ mol})(0.0821)(319 \text{ K})}{1.15 \text{ atm}}$$

8.) (12 pts) What is the partial pressure of nitrogen dioxide if 0.608 mol of nitrogen dioxide is combined with 1.24 mol of oxygen and 0.382 moles of hydrogen gas where the total pressure of the gas is 1.76 atm?

0.608 mol NO2
1.240 mol O2
0.382 mol H2

7.5t

No2
1.76 atm

0.608 mol NO2
1.240 mol O2
0.382 mol H2

1.76 atm

0.479 atm

Please use 0.0821 $^{\text{L}\cdot\text{atm}}/_{\text{mol}\cdot\text{K}}$ as the universal gas constant.